

Inpatient Diabetes Care



INFORMATION FOR HCPs DURING THIS CURRENT CORONAVIRUS PANDEMIC



MONITORING



Check Continuous Blood Glucose monitoring (CBG)

↑ 15 mmol/l x 2 readings

↓ 6 mmol/l x 3 readings

Check capillary ketones **REGARDLESS** of CBG

↑ 1.5 mmol/l



Refer to physician or inpatient DSN team

MEDICATION



STOP

Metformin

SGLT2 inhibitors:

- Empagliflozin (Jardiance)
- Dapagliflozin (Forxiga)
- Canagliflozin (Invokana)
- Ertugliflozin (Steglatro)
- Sotagliflozin (Zynquist)



Until clinician has reviewed

INSULIN



DO NOT STOP INSULIN FOR PEOPLE WITH TYPE 1 DIABETES

REMEMBER

6R's:

- ✓ Right person
- ✓ Right time
- ✓ Right strength
- ✓ Right insulin
- ✓ Right dose
- ✓ Right device

For more information go to ITS training and support: www.inpatientdiabetes.org.uk

For further information contact:

Tel: 0116 258 4674
www.edendiabetes.com

✉ EDEN@uhl-tr.nhs.uk
🐦 @EDEN_Leicester




**INPATIENT DIABETES
TRAINING &
SUPPORT**